

When can I start?

Light work (no heavy lifting) – 2-4 weeks

Heavy work (or above shoulder level) – 12 weeks+

Driving – Discuss with your consultant

Please bring a vest top to all of your follow up physiotherapy appointments.

Your follow up Physiotherapy appointment:

DATE :.....TIME :.....

PHYSIOTHERAPIST :.....

If you have any questions regarding any of the advice included in this leaflet please do not hesitate to contact the physiotherapy department on **0114 263 2112**

For advice on how long to wear your stockings please contact the nursing staff on **0114 263 2176**

Useful website: www.shoulderdoc.co.uk



Claremont Private Hospital
401 Sandygate Road
Sheffield S10 5UB

0114 263 0330
info@claremont-hospital.co.uk
claremont-hospital.co.uk

A Division of Aspen Healthcare Ltd.
Centurion House, 37 Jewry Street London EC3N 2ER
Registration No. 03471084 ©Aspen Healthcare Ltd 2019



issue Feb 2020 / review Feb 2022 / version 1



ACJ Stablisation

Physiotherapy Advice



Welcome to exceptional healthcare



PART OF ASPEN HEALTHCARE

What have I had done?

An operation to stabilise your Acromio-Clavicular Joint (ACJ). There are various techniques available to achieve the desired stability, this will be explained in finer detail by your consultant.

You will be required to wear a sling for a period of time after the operation, this length of time will be decided by your consultant.

Will I experience pain?

You are likely to experience pain in the shoulder and in the upper arm in the first 12 weeks. The pain may be worsened on activity, which is normal. You will be prescribed pain relief to take home, this is to be used to ensure you can meet all recovery goals.

If you find your medication is not managing your pain well, it is important to visit your GP to review the pain relief to avoid falling behind schedule with your rehabilitation.

What should I avoid?

- Removing the sling while arm is still numb
- Moving your arm across your body horizontally
- Heavy lifting in the first 12 weeks.

What can I do to help myself?

Take pain relief as advised by nursing staff.

- Apply ice in a damp pillow case/tea towel to the shoulder for 15 minutes max (before, during or after exercises), ideally 3-5 times a day
- Sleep on your back and use pillows to support the arm for comfort.

Exercises

Never force a stretch, only go to what you can comfortably tolerate. But be progressive.

The exercises will be discussed/demonstrated on the day of your operation. The exercises are very important to allow optimal recovery from your operation.

Start your exercises when the nerve block wears off.



Exercises

Lie on back, use good arm to lift operated arm straight over your head.

Maintain good posture.

Hold 3 seconds.

Repeat 10 times. 4 x daily

Stand facing a wall. Using a towel or paint roller, ease your hand up the wall avoiding compensation (arching the back or shrugging the shoulder). Step closer to the wall to get further with your arm. Reverse down the same way. *Hold 3 seconds, 10 times.* This movement can also be done for the arm out to the side movement.

Sitting or standing. Place rolled up towel/pillow between elbow and body. Maintain 90 degrees at the elbow. Push hand out to the side with a stick, maintain contact between elbow and towel/pillow. Hold 3 seconds.

Repeat 10 times. 4 x daily

Throughout the day, gently move the head and neck. Keeping shoulders still: look up to the ceiling, down to the floor, over left shoulder, over right shoulder, left ear to left shoulder and right ear to right shoulder.